



**THE HONG KONG COLLEGE OF OBSTETRICIANS AND GYNAECOLOGISTS**  
**AND**  
**THE HONG KONG SOCIETY FOR COLPOSCOPY AND CERVICAL PATHOLOGY**



**COA-3 (2021-12)**

### Colposcopy Service Centre Application Form

Name of Applicant	
Position	
Qualification	
Corresponding Address	
E-mail Address	
Phone Number	

Institution/Clinic providing Colposcopy Service	
---	--

**Colposcopists**

	Name	Accreditation cycle (s) e.g. Jan 21 – Dec 23, Jan 24 – Dec 27
1	(Lead Clinician)	
2		
3		
4		
5		
6		

**Sites of colposcopy clinic(s) in the colposcopy service**

1	
2	
3	

**Number clinic(s) in the colposcopy service**

	Mon	Tue	Wed	Thur	Fri	Sat	Sun

Use the number to represent the clinic listed above, specify the nature (D=diagnostic; T=treatment, B=both) and duration of the session

Example: 2, D, 2-5pm



THE HONG KONG COLLEGE OF OBSTETRICIANS AND GYNAECOLOGISTS  
AND  
THE HONG KONG SOCIETY FOR COLPOSCOPY AND CERVICAL PATHOLOGY



## Colposcopy Services

Leading Clinician: \_\_\_\_\_ Signature: \_\_\_\_\_

Period of audit: from \_\_\_\_\_ to \_\_\_\_\_

### Notes on using this Form

1. New application can be made by submitting this application form together with an audit report of colposcopy services. The audit report should cover a 12-month period that falls within the 18 months prior to the date of application.
2. Applying centers can use the annual audit report downloaded from CMS, or use the audit form attached to this form, for reporting of the statistics.
3. At least 50 cases per year are required to apply as Colposcopy Service Centre (COA-08). Therefore, the table on the correlation between colposcopy diagnosis and the worst histology should contain at least 50% of the total number of colposcopy examination submitted, and should have at least 50 cases in total.
4. Applying centers need to fill in the sensitivity and positive predictive value for HSIL, and sensitivity of invasive lesions, by colposcopic examinations performed during the audit period.  
The numbers that are required for the reporting (A – D) can be found in Table 2.
5. At least 60% correlation between colposcopic diagnosis and histological diagnosis of high-grade lesion is expected i.e. sensitivity.
6. 90% of patient treated is expected to have CIN lesion on histological examination.
7. 80% of treatment of CIN is expected to be performed under LA.



THE HONG KONG COLLEGE OF OBSTETRICIANS AND GYNAECOLOGISTS  
AND  
THE HONG KONG SOCIETY FOR COLPOSCOPY AND CERVICAL PATHOLOGY



**Audit Report - Reporting year \_\_\_\_\_ to \_\_\_\_\_**

*(This form can be replaced by the audit report generated from the CMS in HA units)*

Number of colposcopy examinations for patients with new indications \_\_\_\_\_

Number of colposcopy examinations for patients for follow-up \_\_\_\_\_

**Number of Referrals with Cytological Abnormalities**

ASCH	
AGCNOS	
AGC-neoplasia	
HSIL	
AIS	
SCC	
Adenocarcinoma	

**Waiting Time (weeks)**

Suspected carcinoma	
High grade cytological abnormalities	
Low grade cytological abnormalities	



**THE HONG KONG COLLEGE OF OBSTETRICIANS AND GYNAECOLOGISTS**  
**AND**  
**THE HONG KONG SOCIETY FOR COLPOSCOPY AND CERVICAL PATHOLOGY**



**Table 1 - Correlation between Colposcopy Assessment and Histology of Biopsy**

		<b>Histology of Biopsy</b>							
		NORMAL	LG / HPV	HG	MICROIN	INVAS	NO_BX	UNSAT	TOTAL
<b>Colposcopy Diagnosis</b>	NORMAL								
	LG								
	HG								
	MICROIN								
	INVAS								
	UNSAT								
	TOTAL								



**THE HONG KONG COLLEGE OF OBSTETRICIANS AND GYNAECOLOGISTS**  
**AND**  
**THE HONG KONG SOCIETY FOR COLPOSCOPY AND CERVICAL PATHOLOGY**



**Table 2 - Correlation between Colposcopy Assessment and Worst Histology**

		<b>Worst Histology</b>							
		NORMAL	LG / HPV	HG	MICROIN	INVAS	NO_BX	UNSAT	TOTAL
<b>Colposcopy Diagnosis</b>	NORMAL								
	LG								
	HG			(A)					(D)
	MICROIN			(A)					(D)
	INVAS			(A)					(D)
	UNSAT			(C)					
	TOTAL			(B)					

**Sensitivity of colposcopy**

Sensitivity for HSIL or above =  $\frac{\text{No. of HSIL+ by colposcopic examination, (A)}}{\text{Total number of HSIL, (B) - (C)}} \times 100\% = \underline{\hspace{2cm}}$

Positive predictive value for HSIL or above =  $\frac{\text{No. of HSIL+ by colposcopic examination, (A)}}{\text{Total number of colposcopic HSIL+, (D)}} \times 100\% = \underline{\hspace{2cm}}$



THE HONG KONG COLLEGE OF OBSTETRICIANS AND GYNAECOLOGISTS  
AND  
THE HONG KONG SOCIETY FOR COLPOSCOPY AND CERVICAL PATHOLOGY



**Table 3 – Treatments and complications**

	No.	Percentage (%)
Total number of treatment for CIN		NA
Total number of loop excision performed (T)		NA
Number of treatment by LEEP for CIN performed under LA		Out of (T)
Number of other procedures performed for CIN (please specific type of treatment and anaesthesia)		NA
Number of LEEP have histological evidence of CIN		Out of (T)
<b>Number of complications after loop excision:</b>		
Primary haemorrhage		Out of (T)
Secondary haemorrhage		Out of (T)
Infection		Out of (T)
Hospital admission related to procedure		Out of (T)
Others		Out of (T)
Number of loop excision with histological confirmation of CIN		Out of (T)